

Microlocal Analysis of generalized pullbacks of Colombeau functions

Simon Haller, University of Vienna, Faculty of Mathematics

February 01, 2007

Abstract

In distribution theory the pullback of a general distribution by a C^∞ -function is well-defined whenever the normal bundle of the C^∞ -function does not intersect the wavefront set of the distribution. However, the Colombeau theory of generalized functions allows for a pullback by an arbitrary c-bounded generalized function. It has been shown in previous work that in the case of multiplication of Colombeau functions (which is a special case of a C^∞ pullback), the generalized wave front set of the product satisfies the same inclusion relation as in the distributional case, if the factors have their wavefront sets in favorable position. We prove a microlocal inclusion relation for the generalized pullback (by a c-bounded generalized map) of Colombeau functions. The proof of this result relies on a stationary phase theorem for generalized phase functions, which is given in the Appendix. Furthermore we study an example (due to Hurd and Sattinger), where the pullback function stems from the generalized characteristic flow of a partial differential equation.

1 Introduction

The pullback of a general distribution by a C^∞ -function in classical distribution theory, as defined in [7, Theorem 8.2.4], exists if the normal bundle of the C^∞ function intersected with the wavefront set of the distribution is empty. These microlocal restrictions reflect also the well-known fact that in general distribution theory one cannot carry out multiplications unrestrictedly, since the product of two distributions can formally be written as the pullback of a tensor product of the two factors by the diagonal map $\delta : x \mapsto (x, x)$.

Generalized functions in the sense of Colombeau extend distribution theory in a way that it becomes a differential algebra with a product that preserves the classical product $\cdot : C^\infty \times C^\infty \rightarrow C^\infty$. In addition [6, Proposition 1.2.8] states that the Colombeau algebra of generalized functions allows the definition of a pullback by any c-bounded generalized function. The classical concept of a wavefront set has been extended to generalized functions of Colombeau type in [3, 4, 8].

In [10] the microlocal properties of a product of generalized functions were investigated (which can be interpreted as the pullback of a generalized function by a C^∞ function). It was shown that the classical microlocal inclusion relations for the wavefront set of a product of distributions (as in [7, Theorem 8.2.10]) can be extended to generalized functions, if the wavefront sets of the factors are in favorable position.

Furthermore the authors provided a counterexample for a product of Colombeau functions with wavefront sets in unfavorable position that fails the classical microlocal inclusion.

In the present paper we study the wave front set of a generalized pullback of a Colombeau function.

We have divided the paper into four sections:

- In the first section we introduce the concept of a generalized graph.
- In Section 2 we define the transformation of a wave front set by a c-bounded generalized map and the normal bundle of a c-bounded generalized map using the topological concept introduced in the first section.
- The main theorem is stated in the third section. It is a generalization of [7, Theorem 8.2.4].
- Section 4 provides two examples: The counterexample mentioned above, from [10], is investigated in the light of our main theorem. We also study the Hurd-Sattinger example (a partial differential equation of first order with non-smooth coefficient) given in [12], which was further investigated and solved in the Colombeau algebra (it is not solvable in L^1_{loc} , when distributional products are employed) in [9]. The microlocal properties of the generalized solution can be determined using our main theorem, since the solution is the pullback of the initial data (a generalized function) by the characteristic flow (a c-bounded generalized map).
- In the Appendix a stationary phase theorem for generalized phase functions is presented, which is crucial for the proof of our main theorem in section 3.

1.1 Notation and basic notions from Colombeau theory

We use [1, 2, 6, 14] as the standard references for the foundations of Colombeau theory. In the present paper we are working with special Colombeau algebras, denoted by $\mathcal{G}^s$ in [6], although here we will drop the superscript 's' to avoid notational overload.

The Colombeau Algebra: Let us recall the basic construction: A Colombeau (generalized) function on some open set $\Omega \subseteq \mathbb{R}^n$ is defined as equivalence class $(u_\varepsilon)_\varepsilon$ of nets of smooth functions $u_\varepsilon \in C^\infty(\Omega)$ and $\varepsilon \in]0, 1]$ subjected to some asymptotic norm conditions (with respect to ε) for their derivatives on compact sets. We have the following:

- (i) Moderate nets $\mathcal{E}_M(\Omega)$: $(u_\varepsilon)_\varepsilon \in C^\infty(\Omega)^{[0,1]}$ such that, for all $K \subset\subset \Omega$ and $\alpha \in \mathbb{N}^n$, there exists $p \in \mathbb{R}$ such that

$$\sup_{x \in K} \|\partial^\alpha u_\varepsilon(x)\| = O(\varepsilon^{-p}) \text{ as } \varepsilon \rightarrow 0. \quad (1)$$

- (ii) Negligible nets $\mathcal{N}(\Omega)$: $(u_\varepsilon)_\varepsilon \in C^\infty(\Omega)^{[0,1]}$ such that, for all $K \subset\subset \Omega$ and all $q \in \mathbb{R}$ an estimate

$$\sup_{x \in K} \|u_\varepsilon(x)\| = O(\varepsilon^q) \text{ as } \varepsilon \rightarrow 0.$$

holds.

- (iii) $\mathcal{E}_M(\Omega)$ is a differential algebra with operations defined at fixed ε , $\mathcal{N}(\Omega)$ is an ideal and $\mathcal{G}(\Omega) := \mathcal{E}_M(\Omega) / \mathcal{N}(\Omega)$ is the special Colombeau algebra.
- (iv) If we replace the nets of smooth functions by nets of real numbers in (i)-(iii) we obtain the ring of generalized numbers $\widetilde{\mathbb{R}}$.
- (v) There are embeddings, $\sigma : C^\infty(\Omega) \hookrightarrow \mathcal{G}(\Omega)$ as subalgebra and $\iota : \mathcal{D}'(\Omega) \hookrightarrow \mathcal{G}(\Omega)$ as linear space, commuting with partial derivatives.
- (vi) $\Omega \rightarrow \mathcal{G}(\Omega)$ is a fine sheaf and $\mathcal{G}_c(\Omega)$ denotes the subalgebra of elements with compact support; using a cut-off in a neighborhood of the support, one can always obtain representing nets with supports contained in a joint compact set.

Regular Colombeau functions: The subalgebra $\mathcal{G}^\infty$ of regular Colombeau (generalized) functions consists of those elements in $\mathcal{G}(\Omega)$ possessing representatives such that the estimate (1) holds for a certain m uniformly over all $\alpha \in \mathbb{N}^n$.

Rapidly decreasing Colombeau functions: A Colombeau function in $\mathcal{G}(\mathbb{R}^n)$ is called rapidly decreasing in the directions $\Gamma \subseteq S^{n-1}$, if it has a representative with the property that there exists a $N \in \mathbb{N}_0$, such that for all $p \in \mathbb{N}_0$

$$\sup_{(\lambda, \xi_1) \in \mathbb{R}^+ \times \Gamma} (1 + \lambda^2)^{p/2} |u_\varepsilon(\lambda \xi_1)| = O(\varepsilon^{-N}) \text{ as } \varepsilon \rightarrow 0$$

holds.

Fouriertransform of a compactly supported Colombeau function: The Fouriertransform of a $u \in \mathcal{G}_c(\Omega)$ is a well-defined Colombeau function in $\mathcal{G}(\mathbb{R}^n)$ by

$$\mathcal{F}(u) := \left(\int_{\Omega} u_\varepsilon(x) e^{-i\langle x, \cdot \rangle} dx \right)_\varepsilon + \mathcal{N}(\mathbb{R}^n),$$

where $(u_\varepsilon)_\varepsilon$ is a representative of u with joint compact support in Ω .

$\mathcal{G}^\infty$ wavefront set of a Colombeau function: If $v \in \mathcal{G}_c(\Omega)$, we define the set $\Sigma(v) \subset S^{n-1}$ to be the complement of those points having open neighborhoods $\Gamma \subseteq S^{n-1}$ such that $\mathcal{F}(v)$ is rapidly decreasing in the directions Γ . $\Sigma(v)$ is a closed subset of S^{n-1} . Now let $u \in \mathcal{G}(\Omega)$. Then we define the cone of irregular directions at x_0 by

$$\Sigma_{x_0}(u) = \bigcap_{\varphi \in C_c^\infty(\Omega), \varphi(x_0) \neq 0} \Sigma(\varphi \cdot u).$$

Then the (generalized) wave front set of u is the set

$$\text{WF}(u) := \{(x, \xi) \in \Omega \times S^{n-1} \mid \xi \in \Sigma_x(u)\}$$

We denote the projection onto the first component by

$$\text{sing supp}(u) := \{x \in \Omega \mid (x, \xi) \in \text{WF}(u)\}$$

and call this (generalized) singular support of u .

2 The generalized graph

In the first section we introduce the concept of a generalized graph for a c-bounded generalized map. The generalized graph extends the classical graph of a continuous map, in the sense that the generalized graph of the embedded map (which is a c-bounded generalized map) coincides with the classical graph. Furthermore the generalized graph is closed in the product topology of the domain and the image space of the generalized map. This concept will enable us to define the normal bundle of a generalized c-bounded map and the transformation of a wave front set by a generalized c-bounded map (this is carried out in Section 3). Throughout this section we let Ω_1, Ω_2 be open subsets of $\mathbb{R}^n$ resp. $\mathbb{R}^m$.

Definition 2.1. By $\mathcal{G}[\Omega_1, \Omega_2]$ we denote the generalized maps $F \in \mathcal{G}(\Omega_1)^m$ with the property that F is c-bounded on Ω_1 (into Ω_2), i.e. it possesses a representative $(F_\varepsilon)_\varepsilon$ satisfying the condition

$$\forall K \subset\subset \Omega_1 : \exists K' \subset\subset \Omega_2, \exists \varepsilon_0 > 0 : \text{such that } \forall \varepsilon \leq \varepsilon_0 : F_\varepsilon(K) \subseteq K'.$$

Definition 2.2. For $(\lambda_\varepsilon)_\varepsilon$ a net with $\lambda_\varepsilon \in K \subset\subset \Omega_2$, we introduce

$$\text{CP}((\lambda_\varepsilon)_\varepsilon)$$

to denote the set of its clusterpoints.

Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c-bounded generalized map. Then the set

$$\begin{aligned} C_F := & \left\{ (x, y) \in \Omega_1 \times \Omega_2 \mid \exists \text{ a net } (x_\varepsilon)_\varepsilon \text{ in } \Omega_1 : \lim_{\varepsilon \rightarrow 0} x_\varepsilon = x \in \Omega_1 \right. \\ & \text{and } y \in \text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon) \text{ for some map} \\ & \left. \tau :]0, 1] \mapsto]0, 1], \tau(\varepsilon) \leq \varepsilon \right\} \end{aligned}$$

is called the generalized graph of F . Note that we do not consider nets $(x_\varepsilon)_\varepsilon$ in Ω_1 that converge to the boundary $\partial\Omega_1$.

Definition 2.3. Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. F is said to be equi-continuous at x_0 , if there exists a representative (F_ε) which is equi-continuous in x_0 : For all $\gamma > 0$, there exists some $\delta > 0$ such that

$$|F_\varepsilon(x) - F_\varepsilon(x_0)| < \gamma$$

holds for all $x \in B_\delta(x_0)$ and $\varepsilon \in]0, 1]$. The generalized map F is called locally equi-continuous on some subset $X \subseteq \Omega_1$, if it has a representative which is equi-continuous in each point $x \in X$.

Example 2.4. Consider the c -bounded Colombeau function F defined by the representative $(F_\varepsilon)_\varepsilon$

$$F_\varepsilon := \sin(x/\varepsilon),$$

then it is easy to verify by the definition above, that the generalized graph is

$$C_F := \mathbb{R} \times [-1, 1].$$

Example 2.5. Consider the Colombeau function $F := \iota(H)$, where H is the Heavyside function. If we set $g := H * \rho$, where ρ is the mollifier of the embedding ι with $\int \rho(y)dy = 1$, we have that $F_\varepsilon(x) := g(x/\varepsilon)$ defines a representative of F . For all nets $(x_\varepsilon)_\varepsilon$ tending to $x_0 \neq 0$, we obtain that $F_\varepsilon(x_\varepsilon) \rightarrow \text{sign}(x_0)$ as $\varepsilon \rightarrow 0$. Let us consider the nets $(x_\varepsilon)_\varepsilon$ tending to 0: Since we can find zero-nets $(x_\varepsilon)_\varepsilon$ such that $(x_\varepsilon/\varepsilon)$ tends to any point in $\mathbb{R} \cup \{\pm\infty\}$ and g is continuous and bounded, we have that

$$C_F := \{x \in \mathbb{R} \mid x < 0\} \times \{0\} \cup \{0\} \times \overline{\text{Im}(g)} \times \{x \in \mathbb{R} \mid x > 0\} \times \{1\}.$$

Note that $\overline{\text{Im}(g)}$ is a closed interval that contains $[0, 1]$.

Proposition 2.6. Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. If F is locally equi-continuous on Ω_1 , then it follows that the generalized graph of F can be determined pointwise by

$$C_F = \{(x, y) \in \Omega_1 \times \Omega_2 \mid y \in \text{CP}((F_\varepsilon(x))_\varepsilon)\}$$

Proof. We can choose a locally equi-continuous representative $(F_\varepsilon)_\varepsilon$ and proceed along a standard argument: Let $(x_\varepsilon)_\varepsilon, (x'_\varepsilon)_\varepsilon$ be two nets tending to $x_0 \in \Omega_1$. Then we have, using the local equi-continuity of $(F_\varepsilon)_\varepsilon$ on Ω_1 , that for any map $\tau :]0, 1] \rightarrow]0, 1]$ with $\tau(\varepsilon) < \varepsilon$ and all $\gamma > 0$ we can find some $\delta > 0$, such that the distance is bounded by

$$|F_{\tau(\varepsilon)}(x_\varepsilon) - F_{\tau(\varepsilon)}(x'_\varepsilon)| < \gamma$$

for all $x_\varepsilon, x'_\varepsilon \in B_{\delta/2}(x_0)$. Since $\lim_{\varepsilon \rightarrow 0} x_\varepsilon = \lim_{\varepsilon \rightarrow 0} x'_\varepsilon = x_0$ there exists $\varepsilon' \in]0, 1]$ such that $|x_\varepsilon - x'_\varepsilon| < \delta$ holds for all $\varepsilon < \varepsilon'$. Thus $\lim_{\varepsilon \rightarrow 0} F_{\tau(\varepsilon)}(x_\varepsilon) = \lim_{\varepsilon \rightarrow 0} F_{\tau(\varepsilon)}(x'_\varepsilon)$ and it suffices to consider the constant net $x_\varepsilon := x_0$. For any map $\tau :]0, 1] \rightarrow]0, 1]$ with $\tau(\varepsilon) < \varepsilon$ we have that $(F_{\tau(\varepsilon)}(x_0))_\varepsilon$ is a subnet of $(F_\varepsilon(x_0))_\varepsilon$, hence (2.6) follows. $\square$

Lemma 2.7. *Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. If $(x_0, y_0) \in (\Omega_1 \times \Omega_2) \setminus C_F$, then there exist neighborhoods $Y_0 \subseteq \Omega_2$ of y_0 , $X_0 \subseteq \Omega_1$ of x_0 and $\varepsilon' > 0$ such that*

$$Y_0 \cap F_\varepsilon(X_0) = \emptyset$$

for all $\varepsilon < \varepsilon'$.

Proof. Let $(x_0, y_0) \in (\Omega_1 \times \Omega_2) \setminus C_F$. The proof proceeds by contradiction: Assume that for all neighborhoods $Y' \subseteq \Omega_2$ of y_0 and $X' \subseteq \Omega_1$ of x_0 and for all $\varepsilon' > 0$ there exists some $\tau = \tau(X', Y', \varepsilon') < \varepsilon'$ with

$$Y' \cap F_{\tau(\varepsilon)}(X') \neq \emptyset.$$

Then for all $\varepsilon \in]0, 1]$, by setting $X' := B_\varepsilon(x_0) \cap \Omega_1$, $Y' := B_\varepsilon(y_0) \cap \Omega_2$ and $\varepsilon' := \varepsilon$, we can find some $x_\varepsilon \in B_\varepsilon(x_0) \cap \Omega_1$ and $\tau(\varepsilon) < \varepsilon$ such that $F_{\tau(\varepsilon)}(x_\varepsilon) \in B_\varepsilon(y_0) \cap \Omega_2$. Now $(x_\varepsilon)_\varepsilon$ is a convergent net in Ω_1 with $\lim_{\varepsilon \rightarrow 0} x_\varepsilon = x_0 \in \Omega_1$ and the net $(F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon$ converges to y_0 . Since $\varepsilon \mapsto \tau(\varepsilon)$ defines a map with $\tau(\varepsilon) < \varepsilon$ and $y_0 \in \text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon)$ we have obtained $(x_0, y_0) \in C_F$, which is a contradiction. $\square$

Remark 2.8. *Let X, Y be locally compact topological Hausdorff spaces. Assume that $A \subseteq X \times Y$ is a closed set (in the product topology) with the property that for all compact sets $X_0 \subset\subset X$, it holds that $A \cap (X_0 \times Y)$ is compact. Then it follows that the projection onto X*

$$\begin{aligned} \pi_X : X \times Y &\rightarrow X \\ (x, y) &\mapsto x \end{aligned}$$

has the property that $\pi_X|_A$ is proper and is thus closed. In particular $\pi_X(A)$ is a closed set.

Proposition 2.9. *Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map, then the generalized graph C_F has the following properties:*

- (i) *It extends the classical notion of a graph (of a continuous function) in the following sense: If F is the embedding of a continuous function $G := (G_1, \dots, G_m) \in C(\Omega_1, \Omega_2)$, i.e. $F = (\iota(G_1), \dots, \iota(G_m))$, then the generalized graph of F coincides with the graph of the continuous function G .*
- (ii) *C_F is a closed set in the product topology of $\Omega_1 \times \Omega_2$ and*
- (iii) *for any compact set $X_0 \subset\subset \Omega_1$ it holds that $C_F \cap (X_0 \times \Omega_2)$ is a compact subset of $\Omega_1 \times \Omega_2$.*

Proof. (i) follows by Proposition 2.6 and the fact that the embedding of a continuous function is a locally equi-continuous generalized function.

(ii) Let $(x_0, y_0) \in (\Omega_1 \times \Omega_2) \setminus C_F$, then we know by Lemma 2.7 that there exists $X_0 \subseteq \Omega_1, Y_0 \subseteq \Omega_2$ open neighborhoods of x_0 resp. y_0 and $\varepsilon' > 0$, such that

$$F_\varepsilon(X_0) \cap Y_0 = \emptyset$$

for all $\varepsilon < \varepsilon'$. Now for any point $x \in X_0$ and any net $(x_\varepsilon)_\varepsilon$ in Ω_1 converging to x , there exists some $\varepsilon' > 0$ such that $x_\varepsilon \in X_0$ for all $\varepsilon < \varepsilon'$. It follows that

$$C_F \cap (X_0 \times Y_0) = \emptyset,$$

so $(\Omega_1 \times \Omega_2) \setminus C_F$ is open in $\Omega_1 \times \Omega_2$ and hence C_F is closed in the product topology of $\Omega_1 \times \Omega_2$.

(iii) Since F is c -bounded we can find some $\varepsilon' > 0$ and a compact neighborhood $K \subset\subset \Omega_1$ of X_0 and $K' \subset\subset \Omega_2$, such that

$$F_\varepsilon(K) \subseteq K'$$

for all $\varepsilon < \varepsilon'$. Now if $(x, y) \in C_F \cap (X_0 \times \Omega_2)$, there exists a net $(x_\varepsilon)_\varepsilon$ in Ω_1 converging to x , such that $y \in \text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon)$, where $\tau :]0, 1] \rightarrow]0, 1]$ is some map with $\tau(\varepsilon) < \varepsilon$. Since $(x_\varepsilon)_\varepsilon$ converges to x and K is a neighborhood of x there exists $\varepsilon' > 0$, such that $x_\varepsilon \in K$ for all $\varepsilon < \varepsilon'$. It follows that $(F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon \in K'$ for all $\varepsilon < \varepsilon'$. So we have that $\text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon) \subseteq K'$ and $(x, y) \in K \times K'$. We conclude that $C_F \cap (X_0 \times \Omega_2) = C_F \cap (X_0 \times \Omega_2) \cap K \times K'$ is a compact subset of $\Omega_1 \times \Omega_2$. $\square$

Lemma 2.10. *Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. Furthermore, let $X_0 \subset\subset \Omega_1$ and $Y_0 := \pi_2(C_F \cap (X_0 \times \Omega_2))$. If Y is a neighborhood of Y_0 , then there exists a neighborhood $X \subseteq \Omega_1$ of X_0 and $\varepsilon' > 0$, such that*

$$F_\varepsilon(X) \subseteq Y$$

for all $\varepsilon < \varepsilon'$.

Proof. First we proof the Lemma for $X_0 := \{x_0\}$ containing only a single point $x_0 \in \Omega_1$. We note that Y_0 is a compact subset of Ω_2 by Proposition 2.9. Assume that the statement of the Lemma does not hold, that is for all neighborhoods X' of x_0 and for all $\varepsilon > 0$ there exists some $\tau(\varepsilon) < \varepsilon$ and $x_\varepsilon \in X'$ with

$$F_{\tau(\varepsilon)}(x_\varepsilon) \notin Y.$$

Then (by setting $X' = B_\varepsilon(x_0) \cap \Omega_1$) for all $\varepsilon \in]0, 1]$, we can find a $x_\varepsilon \in B_\varepsilon(x_0) \cap \Omega_1$ and $\tau(\varepsilon)$ with $\tau(\varepsilon) < \varepsilon$ such that $F_{\tau(\varepsilon)}(x_\varepsilon) \notin Y$.

Since Y is a neighborhood of Y_0 , we have that $\overline{Y^c} \cap Y_0 = \emptyset$. The net $(x_\varepsilon)_\varepsilon$ was chosen to converge to x_0 and the net $(F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon \in Y^c$ has the property that the set of clusterpoints $\text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon)$ is contained in the closure of Y^c . By the definition of the generalized graph C_F we have that $\text{CP}((F_{\tau(\varepsilon)}(x_\varepsilon))_\varepsilon) \subseteq Y_0$, which contradicts the fact that $\overline{Y^c} \cap Y_0 = \emptyset$.

Now we consider the general case, when X_0 is an arbitrary compact set. Then Y is a neighborhood of $\pi_2(C_F \cap X_0 \times \Omega_2)$ and it follows that Y is a neighborhood of each set $\pi_2(C_F \cap \{z\} \times \Omega_2)$ for $z \in X_0$. We can apply the first part of the proof for each point z and obtain (open) neighborhoods X_z of

each point z such that $F_\varepsilon(X_z) \subseteq Z$ for all $\varepsilon < \varepsilon_z$, where $\varepsilon_z \in]0, 1]$ depends on z . Since X_0 is compact and $(X_z)_{z \in X_0}$ is an open covering we can choose some finite subcovering $(X_{z_k})_{k=1}^l$. Then $X := \bigcup_{k=1}^l X_{z_k}$ is a neighborhood of X_0 such that $f_\varepsilon(X) \subseteq Y$ holds for $\varepsilon < \varepsilon' := \min_{k=1}^l \varepsilon_{z_k}$. $\square$

Corollary 2.11. *Let $F \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. Furthermore let $X_0 \subset\subset \Omega_1$ and $Y_0 := \pi_2(C_F \cap X_0 \times \Omega_2)$. If $Y \subseteq \Omega_2$ is some neighborhood of Y_0 , then there exists a neighborhood $X \subseteq \Omega_1$ of X_0 , such that*

$$C_F \cap (X \times \Omega_2) \subseteq X \times Y. \quad (2)$$

Proof. If X' is the neighborhood constructed in Lemma 2.10, then we can find some $\delta > 0$, such that $B_\delta(X_0) \subseteq X'$. Set $X := B_{\delta/2}(X_0)$, then the statement is true, since X' is a neighborhood for each point of X . $\square$

3 Transformation of wave-front sets

In this section we consider a c -bounded generalized map $f \in \mathcal{G}[\Omega_1, \Omega_2]$, where $\Omega_1 \subseteq \mathbb{R}^n$, $\Omega_2 \subseteq \mathbb{R}^m$ are open sets and $\Gamma \subseteq \Omega_2 \times S^{m-1}$ is a closed set. We define the normal bundle of a c -bounded generalized map and consider the transformation of wavefront sets by c -bounded generalized maps.

Definition 3.1. *Let $y \in \Omega_2$ and $Y \subset\subset \Omega_2$ be some compact set. Then we use the notation*

$$\Gamma_y := \{\eta \in S^{m-1} \mid (y, \eta) \in \Gamma\}$$

and

$$\Gamma_Y := \{\eta \in S^{m-1} \mid \exists y \in Y : (y, \eta) \in \Gamma\} = \bigcup_{y \in Y} \Gamma_y.$$

Note that both sets are closed subsets of S^{m-1} , since

$$\Gamma_Y = \pi_2(\Gamma \cap (Y \times S^{m-1}))$$

is the image of a compact set under the continuous projection

$$\pi_2 : (y, \eta) \mapsto \eta.$$

Definition 3.2. *According to [11, Definition 2.2] a net $(\lambda_\varepsilon)_\varepsilon$ in $\mathbb{R}$ is said to be of slow-scale, if*

$$\exists \varepsilon' \in]0, 1] : \forall t \geq 0, \exists C_t > 0 \text{ such that } |\lambda_\varepsilon|^t \leq C_t \varepsilon^{-1}, \text{ for all } \varepsilon < \varepsilon'$$

holds.

Lemma 3.3. *Let $\Omega \subseteq \mathbb{R}^m$ be an open set and let Γ be some closed subset of $\Omega \times S^{m-1}$. Furthermore let $Y_0 \subset\subset \Omega$ be a compact set. If $V \subseteq S^{m-1}$ is some closed neighborhood of Γ_{Y_0} , then there exists a neighborhood Y of Y_0 such that*

$$\Gamma_Y \subseteq V$$

holds.

Proof. We note that Γ_{Y_0} is a compact subset of S^{m-1} . For any compact set $L \subset \mathbb{R}^m$ and $\lambda > 0$ we define the set $K_\lambda(L) := \{z \in \mathbb{R}^m \mid \exists z' \in L : |z - z'| \leq \lambda\}$. Since V is a compact neighborhood of the compact set Γ_{Y_0} there exists some $\delta > 0$ such that $K_\delta(\Gamma_{Y_0}) \cap S^{m-1} \cap (S^{m-1} \setminus V) = \emptyset$. We prove the statement by contradiction: Assume the statement is false, then we have that for all $\varepsilon \in]0, 1]$ there exists a point $y_\varepsilon \in K_{\delta \cdot \varepsilon}(Y_0)$ with the property that $\Gamma_{y_\varepsilon} \not\subseteq V$. Therefore we can find some $\xi_\varepsilon \in \Gamma_{y_\varepsilon} \setminus V \neq \emptyset$ for each $\varepsilon \in]0, 1]$. This leads to a net $(y_\varepsilon, \xi_\varepsilon)_\varepsilon$ contained in the compact set $\Gamma \cap (S_\delta(Y_0) \times S^{m-1})$. Thus the net $(y_\varepsilon, \xi_\varepsilon)_\varepsilon$ has a finer net $(y_{\tau(\varepsilon)}, \xi_{\tau(\varepsilon)})_\varepsilon$ that converges to some $(y_0, \xi_0) \in \Gamma \cap (K_\delta(Y_0) \times S^{m-1})$ and $\xi_0 \in \Gamma_{y_0}$. Since $y_\varepsilon \in K_{\delta \cdot \varepsilon}(Y_0)$, it follows that the clusterpoint y_0 is contained in Y_0 . We conclude that $\xi_0 \in \Gamma_{y_0} \subseteq \Gamma_{Y_0}$. We have that there exists $\varepsilon_1 \in]0, 1]$ such that

$$\max(|\xi_{\tau(\varepsilon)} - \xi_0|, |y_{\tau(\varepsilon)} - y_0|) < \delta$$

holds for all $\varepsilon < \varepsilon_1$. We have chosen δ such that $\{\xi \in S^{m-1} \mid \exists \xi_0 \in \Gamma_{Y_0} : |\xi - \xi_0| < \delta\} \subseteq V$. It follows that $\xi_{\tau(\varepsilon)} \in V$ for all $\varepsilon < \varepsilon_1$, which contradicts the choice $\xi_\varepsilon \in \Gamma_{y_\varepsilon} \setminus V$. $\square$

Although the pullback of a Colombeau function u by any c -bounded generalized map f is well-defined, we cannot derive a general microlocal inclusion for the pullback without requiring further properties for the generalized map f . We define an open subdomain D_f of $\Omega_1 \times S^{m-1}$, where the generalized map $(x, \eta) \mapsto {}^T df_\varepsilon(x)\eta$ has certain properties which are needed to obtain a microlocal inclusion relation. This leads to the notion of a generalized normal bundle.

Definition 3.4. Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map, then we define the open set D_f by

$$\begin{aligned} D_f := & \{(x, \eta) \in \Omega_1 \times S^{m-1} \mid \exists \text{ neighborhood } X \times V \subseteq \Omega_1 \times S^{m-1} \text{ of } (x, \eta) \\ & \text{and a positive net of slow growth } (\sigma_\varepsilon)_\varepsilon, \exists \varepsilon' \in]0, 1] : \\ & \inf_{(x, \eta) \in X \times V} |\sigma_\varepsilon {}^T df_\varepsilon(x)\eta| \geq 1 \text{ and} \\ & \sup_{(x, \eta) \in X \times V^\perp} |\sigma_\varepsilon {}^T df_\varepsilon(x)\eta| \leq 1 \text{ for all } \varepsilon < \varepsilon'\}, \end{aligned}$$

where $V^\perp := \{\eta \in S^{m-1} \mid \exists \eta_0 \in V : \langle \eta, \eta_0 \rangle = 0\}$. Then the generalized normal bundle of f is defined by

$$N_f := \{(y, \eta) \in \Omega_2 \times S^{m-1} \mid (x, y) \in C_f, (x, \eta) \notin D_f\}.$$

This is the analog to the classical normal bundle. Furthermore, we define the wave-front unfavorable support of f with respect to a closed set $\Gamma \subseteq \Omega_2 \times S^{m-1}$ by

$$U_f(\Gamma) := \{x \in \mathbb{R}^n \mid (x, y) \in C_f, (x, \eta) \notin D_f, (y, \eta) \in \Gamma\}.$$

Example 3.5. We consider the c -bounded generalized map defined by

$$f_\varepsilon(x, y) = (x + \gamma_\varepsilon y, x - \gamma_\varepsilon y),$$

where $\lim_{\varepsilon \rightarrow 0} \gamma_\varepsilon = 0$.

Then the transposed Jacobian is

$${}^T df_\varepsilon(x, y) := \begin{pmatrix} 1 & 1 \\ \gamma_\varepsilon & -\gamma_\varepsilon \end{pmatrix}$$

and ${}^T df_\varepsilon(x, y)\eta = (\eta_1 + \eta_2, \gamma_\varepsilon(\eta_1 - \eta_2))$. Let $\tilde{\eta} := (\pm 1, \mp 1)$ and $\tilde{\eta}^\perp := (\pm 1, \pm 1)$ then

$$\begin{aligned} a_\varepsilon(x, \tilde{\eta}) &:= |\sigma_\varepsilon {}^T df_\varepsilon(x) \tilde{\eta}| = \gamma_\varepsilon \sigma_\varepsilon 2 \\ b_\varepsilon(x, \tilde{\eta}) &:= |\sigma_\varepsilon {}^T df_\varepsilon(x) \tilde{\eta}^\perp| = \sigma_\varepsilon 2 \end{aligned}$$

shows that for any choice of a slow-scaling net σ_ε either $a_\varepsilon \rightarrow 0, b_\varepsilon \rightarrow 1 \in \mathbb{R}^+$ or $a_\varepsilon \rightarrow 1 \in \mathbb{R}^+, b_\varepsilon \rightarrow \infty$ (the second case is ruled out if γ_ε is not a slow-scaled net). For all other $\eta \notin S^{n-1} \setminus (\pm 1, \mp 1)$ it is easy to find a neighborhood V of η and some slow-scaling net σ_ε , such that $\inf_{\mathbb{R}^2 \times V} a_\varepsilon(x, \eta) \geq 1$ and $\sup_{\mathbb{R}^2 \times V^\perp} a_\varepsilon(x, \eta) \leq 1$ for ε small enough. It follows that $D_f := \mathbb{R}^2 \times S^1 \setminus (\pm 1, \mp 1)$.

Lemma 3.6. *The generalized normal bundle N_f and the wave-front unfavorable support $U_f(\Gamma)$ of f (with respect to Γ) are closed sets. If $N_f \cap \Gamma = \emptyset$, then $U_f(\Gamma) = \emptyset$.*

Proof. Let $x_0 \notin U_f(\Gamma)$, then since D_f is open there exist neighborhoods X_0 resp. V_0 of x_0 resp. Γ_{Y_0} such that $X_0 \times V_0 \subseteq D_f$. By Lemma 3.3 there exists a neighborhood Y of Y_0 , such that V_0 is still a neighborhood of Γ_Y . Corollary 2.11 provides a neighborhood X' of x_0 such that $C_f \cap (X' \times \Omega_2) \subseteq X' \times Y$. Now we can choose a smaller neighborhood $X_1 \subset X' \cap X_0$ (such that $X' \cap X_0$ is a neighborhood of X_1) of x_0 , such that $X_1 \times V_0 \subseteq D_f$ and $C_f \cap (X_1 \times \Omega_2) \subseteq X_1 \times Y$. For all $x_1 \in X_1$ and $y \in \Omega_2$ with $(x_1, y) \in C_f$, it holds that $(x_1, y) \in X_1 \times Y$ and if $(y, \eta) \in \Gamma$ it follows that $\eta \in \Gamma_Y \subseteq V$ and thus $(x_1, \eta) \in D_f$. It follows that $x_1 \notin U_f$ for all $x_1 \in X_1$, $X_1 \cap U_f = \emptyset$ and thus $U_f(\Gamma)^c$ is an open set. $\square$

A substantial step in the proof of the main theorem applies a generalized stationary phase theorem (in the Appendix). In order to obtain a lower bound for the gradient of the occurring phase function we have to consider the map

$$\begin{aligned} \Omega_1 \times S^{m-1} \times S^{n-1} &\rightarrow S^{n-1} \\ (x, \eta, \xi) &\mapsto \left| \frac{{}^T df_\varepsilon(x) \eta}{|{}^T df_\varepsilon(x) \eta|} - \xi \right|. \end{aligned}$$

So we introduce the following notation:

Lemma 3.7. *Consider*

$$M_\varepsilon(x, \eta) := \frac{Tdf_\varepsilon(x)\eta}{|Tdf_\varepsilon(x)\eta|}$$

on the domain D_f , then $\eta \mapsto M_\varepsilon(x, \eta)$ defines an equi-continuous Colombeau function in the η variable at fixed x .

Proof. The map $g : \xi \rightarrow \frac{\xi}{|\xi|}$ is equi-continuous on $\mathbb{R}^n \setminus B_\delta(0)$ for any fixed $\delta > 0$ since

$$\left| \frac{\xi}{|\xi|} - \frac{\eta}{|\eta|} \right| = \frac{1}{|\xi|} \left| \xi - \frac{\eta|\xi|}{|\eta|} \right| = \frac{1}{|\xi|} \left| \xi - \eta + \frac{\eta}{|\eta|} (|\eta| - |\xi|) \right| \leq \frac{2}{|\xi|} |\xi - \eta|.$$

Let $(x_0, \eta_0) \in D_f$, then we can find some neighborhood $X \times V \subseteq \Omega_1 \times S^{m-1}$ of (x_0, η_0) such that

$$\inf_{(x, \eta) \in X \times V} |\sigma_\varepsilon Tdf_\varepsilon(x)\eta| \geq 1 \text{ and } \sup_{(x, \eta) \in X \times V^\perp} |\sigma_\varepsilon Tdf_\varepsilon(x)\eta| \leq 1$$

for some slow-scaling net $(\sigma_\varepsilon)_\varepsilon$. Then we conclude that

$$\begin{aligned} |M_\varepsilon(x, \eta) - M_\varepsilon(x, \xi)| &\leq \frac{2}{|Tdf_\varepsilon(x)\eta|} |Tdf_\varepsilon(x)\eta - Tdf_\varepsilon(x)\xi| \\ &= \frac{2}{|\sigma_\varepsilon Tdf_\varepsilon(x)\eta|} |\sigma_\varepsilon Tdf_\varepsilon(x)(\eta - \xi)| \leq \frac{\sup_{(x, \xi) \in X \times V^\perp} |\sigma_\varepsilon Tdf_\varepsilon(x)\xi|}{\inf_{(x, \eta) \in X \times V} |\sigma_\varepsilon Tdf_\varepsilon(x)\eta|} \\ &\quad \cdot |\xi - \eta| \leq 2|\xi - \eta| \end{aligned}$$

uniformly for all $x \in X_0$ and $\xi, \eta \in V$. $\square$

Remark 3.8. *From the preceding Lemma we obtain that if $\eta_\varepsilon \rightarrow \eta_0$ and $x_\varepsilon \rightarrow x_0$, then*

$$\text{CP}((M_\varepsilon(x_\varepsilon, \eta_\varepsilon))_\varepsilon) = \text{CP}((M_\varepsilon(x_\varepsilon, \eta_0))_\varepsilon),$$

which simplifies the determination of the generalized graph C_M .

Definition 3.9. *Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map and assume that $D_f \neq \emptyset$. We define the transformation map Φ_f by*

$$\begin{aligned} \Phi_f : D_f &\rightarrow \Omega_2 \times S^{n-1} \\ (x, \eta) &\mapsto \left(f_\varepsilon(x), \frac{Tdf_\varepsilon(x)\eta}{|Tdf_\varepsilon(x)\eta|} \right). \end{aligned}$$

Consider the projections

$$\begin{aligned} \Pi_{\Phi_f, 1} : C_{\Phi_f} &\rightarrow \Omega_1 \times S^{n-1}, \quad (x, \eta, y, \xi) \mapsto (x, \xi) \\ \Pi_{\Phi_f, 2} : C_{\Phi_f} &\rightarrow \Omega_2 \times S^{m-1}, \quad (x, \eta, y, \xi) \mapsto (y, \eta) \end{aligned}$$

and define the set $f^*\Gamma$ by

$$\begin{aligned} f^*\Gamma &:= \Pi_{\Phi_f,1} \circ \Pi_{\Phi_f,2}^{-1}(\Gamma) \\ &= \{(x, \xi) \in \Omega_1 \times S^{n-1} \mid (x, \eta, y, \xi) \in C_{\Phi_f}, (y, \eta) \in \Gamma\}. \end{aligned}$$

In the case of $D_f = \emptyset$, we set $f^*\Gamma := \Omega_1 \times S^{n-1}$.

Lemma 3.10. *Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map and let $\Gamma \subset \Omega_2 \times S^{m-1}$ be a closed set in the product topology of $\Omega_2 \times S^{m-1}$. Then the set $f^*\Gamma$ is closed in the product topology of $\Omega_1 \times S^{n-1}$.*

Proof. The maps

$$\begin{aligned} \Pi_1 : \Omega_1 \times S^{m-1} \times \Omega_2 \times S^{n-1} &\rightarrow \Omega_1 \times S^{n-1}, \\ (x, \eta, y, \xi) &\mapsto (x, \xi) \\ \Pi_2 : \Omega_1 \times S^{m-1} \times \Omega_2 \times S^{n-1} &\rightarrow \Omega_2 \times S^{m-1}, \\ (x, \eta, y, \xi) &\mapsto (y, \eta) \end{aligned}$$

are continuous projections and we have that

$$\Pi_{\Phi_f,1} = \Pi_1|_{C_{\Phi_f}}$$

and

$$\Pi_{\Phi_f,2} = \Pi_2|_{C_{\Phi_f}}.$$

C_{Φ_f} has the following property: For all $K \subset \subset \Omega_1 \times S^{n-1}$ there exists some $K' \subset \subset \Omega_1 \times S^{m-1} \times \Omega_2 \times S^{n-1}$ such that $C_{\Phi_f} \cap \Pi_1^{-1}(K) \subseteq K'$ (cf. Proposition 2.9). By Remark 2.8 the map $\Pi_{\Phi_f,1}$ is thus closed. It follows that $\Pi_{\Phi_f,1} \circ \Pi_{\Phi_f,2}^{-1}(\Gamma)$ is a closed set in the subset topology of $\Omega_1 \times S^{n-1}$. $\square$

4 Generalized pullbacks of Colombeau functions

In this section we prove the main result which gives a microlocal inclusion relation for the generalized pullback of a Colombeau function. The proof of the theorem relies on a generalized stationary phase theorem, the details of which are discussed in the Appendix.

Definition 4.1. *Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. Then we call f slow-scaled in all derivatives on the open set $X_0 \subseteq \Omega_1$, if for all $\alpha \in \mathbb{N}^n$ there exists slow-scaled nets $(r_{\alpha,\varepsilon})_\varepsilon$ such that*

$$\sup_{X_0} |\partial^\alpha f_\varepsilon(x)| \leq C_\alpha r_{\alpha,\varepsilon} \text{ as } \varepsilon \rightarrow 0 \quad (3)$$

holds, where C_α are constants. Furthermore we call f slow-scaled in all derivatives at $x_0 \in \Omega_1$, if there exists a neighborhood X_0 of x_0 such that (3) holds for all $\alpha \in \mathbb{N}^n$.

Define the sets

$$S_f := \{x \in \Omega_1 \mid f \text{ is slow - scaled in all derivatives at } x\}$$

and

$$K_f(u) := \bigcap_{k \in \tilde{\mathbb{R}}^m} \pi_1(C_f \cap \text{supp}(u - k) \times \Omega_2).$$

Remark 4.2. The sets $\pi_1(C_f \cap \Omega_1 \times \text{supp}(u - k))$ are closed because $\pi_1|_{C_f}$ is a proper map and the sets $\text{supp}(u - k) \subset \Omega_1$ and $C_f \subseteq \Omega_1 \times \Omega_2$ are closed. Thus the set $K_f(u)$ is closed in the relative topology of $\Omega_1 \subseteq \mathbb{R}^n$.

Lemma 4.3. Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map. Furthermore let $x_0 \notin U_f(\Gamma) \cup S_f^c$ and $Y_0 := \pi_2(C_f \cap \{x_0\} \times \Omega_2)$. If W is some open neighborhood of $(f^*\Gamma)_{x_0}$, then there exist neighborhoods X of x_0 , V of Γ_{Y_0} and Y of Y_0 with the following properties:

$$\begin{aligned} X \times V &\subseteq D_f(\Gamma), \quad (f^*\Gamma)_X \subseteq W, \quad f_\varepsilon(X) \subseteq Y \text{ for all } \varepsilon < \varepsilon', \\ (f_\varepsilon) &\text{ is slow - scaling in all derivatives on } X \text{ and } \Gamma_Y \subseteq V \end{aligned}$$

Furthermore there exists a positive constant $c > 0$, such that

$$\inf_{(x, \eta, \xi) \in X \times V \times W^c} |M_\varepsilon(x, \eta) - \xi| > c \text{ for all } \varepsilon < \varepsilon'$$

holds (we are using the notation from Lemma 3.7).

Proof. From $x_0 \notin U_f(\Gamma)$ it follows that $\{x_0\} \times \Gamma_{Y_0}$ is a compact subset of the open set D_f . Thus we can find some neighborhood $X' \times V' \subseteq D_f$ of $\{x_0\} \times \Gamma_{Y_0}$. Since $x_0 \in S_f$ we can assume without loss of generality that f is slow-scale in all derivatives on the compact set X' .

Since W is a neighborhood of $(f^*\Gamma)_{x_0}$ we have that

$$(f^*\Gamma)_{x_0} = \pi_2(C_M \cap \{x_0\} \times \Gamma_{Y_0} \times S^{m-1}) \subset\subset W$$

where C_M denotes the generalized graph of the generalized map defined by $(x, \eta) \mapsto M_\varepsilon(x, \eta)$ on the open domain D_f . By Lemma 2.10 there exist neighborhoods X'', V'' of x_0 resp. Γ_{Y_0} such that

$$M_\varepsilon(X'' \times V'') \subseteq W \tag{4}$$

for all $\varepsilon < \varepsilon'$. Let $V := V' \cap V''$, then by Lemma 3.3 we can find some neighborhood Y of Y_0 such that $\Gamma_Y \subseteq V$. By Lemma 2.10 there exists some neighborhood X''' of x_0 such that $f_\varepsilon(X''') \subseteq Y$ for small ε . Let $X := X' \cap X'' \cap X'''$ and $Z := X \times V \times W^c$. Then there exists $(x_\varepsilon, y_\varepsilon, \xi_\varepsilon) \in Z$ (note that $(x, \eta, \xi) \mapsto |M_\varepsilon(x, \eta) - \xi|$ is a continuous function for each $\varepsilon \in]0, 1]$ and Z is a compact set) such that

$$c_\varepsilon := \inf_{(x, \eta, \xi) \in Z} |M_\varepsilon(x, \eta) - \xi| = |M_\varepsilon(x_\varepsilon, \eta_\varepsilon) - \xi_\varepsilon|$$

holds for some net $(x_\varepsilon, \eta_\varepsilon, \xi_\varepsilon)_\varepsilon$. By (4) we have that $c_\varepsilon > c > 0$ holds for all $\varepsilon < \varepsilon'$, where c is a constant. $\square$

Theorem 4.4. *Let $f \in \mathcal{G}[\Omega_1, \Omega_2]$ be a c -bounded generalized map with representative $(f_\varepsilon)_\varepsilon$. For $u \in \mathcal{G}(\Omega_2)$ with representative $(u_\varepsilon)_\varepsilon$, we define the pullback f^*u by*

$$f^*u := (u_\varepsilon(f_\varepsilon(x)))_\varepsilon + \mathcal{N}(\Omega_1).$$

*Then f^*u is well-defined and the microlocal inclusion relation*

$$\text{WF}(f^*u) \subseteq f^*\text{WF}(u) \bigcup (U_f(\text{WF}(u)) \times S^{n-1}) \bigcup ((K_f(u) \cap S_f^c) \times S^{n-1})$$

holds.

Proof. According to [6, Proposition 1.2.8] the pullback f^*u is a well-defined Colombeau function in $\mathcal{G}(\Omega_1)$.

Let $x_0 \in \Omega_1$ and $Y_0 = \{y \in \Omega_2 \mid (x_0, y) \in C_f(\Omega_1)\}$ and ξ_0 . We set $\Gamma := \text{WF}(u)$ and $\Gamma_y := \{\eta \mid (y, \eta) \in \Gamma\}$.

Suppose that $(x_0, \xi_0) \notin f^*\Gamma \bigcup U_f(\Gamma) \times S^{n-1} \bigcup (K_f(u) \cap S_f^c) \times S^{n-1}$, so $\xi_0 \notin (f^*\Gamma)_{x_0}$ and $x_0 \notin U_f(\Gamma) \bigcup K_f(u) \cap S_f^c$. We distinguish the cases $x_0 \notin K_f(u) \cup U_f(\Gamma)$ and $x_0 \notin S_f^c \cup U_f(\Gamma)$:

In the first case we have that there exists a generalized constant $k \in \widetilde{\mathbb{R}}^n$ with representative $(k_\varepsilon)_\varepsilon$, such that $C_f \cap \{x_0\} \times \text{supp}(u - k) = \emptyset$. Hence for all $y \in Y_0$ it holds that $y \notin \text{supp}(u - k)$. Since Y_0 and $\text{supp}(u - k)$ are two disjoint closed sets, we can find a closed neighborhood Y of Y_0 such that $Y \cap \text{supp}(u) = \emptyset$. Let $\chi \in C_c^\infty(\Omega_2)$ with the property that $\chi \equiv 1$ on some compact neighborhood $Y' \subseteq Y^\circ$ of Y_0 and $\text{supp}(\chi) \subseteq Y$. By Lemma 2.10 there exists a neighborhood X' of x_0 and $\varepsilon' \in]0, 1]$ such that $f_\varepsilon(X') \subseteq Y'$ for $\varepsilon < \varepsilon'$.

In order to show that $\xi_0 \notin (\text{WF}(f^*u))_{x_0}$ for $x_0 \notin K_f(u)$ we have to find a smooth function φ with support on a neighborhood of x_0 , such that $\mathcal{F}(f^*u\varphi)$ is rapidly decreasing on some neighborhood of ξ_0 .

We choose φ to be a smooth function with $\text{supp}(\varphi) \subseteq X'$. For all $\varepsilon < \varepsilon'$ the identity $f_\varepsilon^*u \cdot \varphi = f_\varepsilon^*(\chi u) \cdot \varphi$ holds, since the functions χ and φ were chosen such that $(\chi \circ f_\varepsilon) \cdot \varphi \equiv 1$ for $\varepsilon < \varepsilon'$. We have that $\chi \cdot (u_\varepsilon - k_\varepsilon) \in \mathcal{N}(\Omega_2)$ and it follows that $f^*u \cdot \varphi$ is $\mathcal{G}^\infty$, so $\xi_0 \notin (\text{WF}(f^*u))_{x_0}$.

Now we consider the case $x_0 \notin U_f(\Gamma) \cup S_f^c$. Let W be an open set in S^{n-1} such that W^c is a neighborhood of ξ_0 . Lemma 4.3 implies that there exist neighborhoods X of x_0 , Y of Y_0 and V of Γ_{Y_0} such that $(f^*\Gamma)_X \subseteq W$, $f_\varepsilon(X) \subseteq Y$ for all $\varepsilon < \varepsilon'$, $\Gamma_Y \subseteq V$, (f_ε) is slow-scaling in all derivatives on X and $\inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} |M_\varepsilon(x, \eta_1) - \xi_1| > d > 0$ holds for all $\varepsilon < \varepsilon'$, where d is some positive constant.

Let $\chi \in C_c^\infty(\Omega_2)$ with the property that $\chi \equiv 1$ on some compact neighborhood $Y' \subseteq Y^\circ$ of Y_0 and $\text{supp}(\chi) \subseteq Y$. By Lemma 2.10 there exists a neighborhood X' of x_0 and $\varepsilon' \in]0, 1]$ such that $f_\varepsilon(X') \subseteq Y'$ for $\varepsilon < \varepsilon'$. Without loss of generality we can assume that $X' \subseteq X$.

In order to show that $\xi_0 \notin \text{WF}(f^*u)_{x_0}$ for $x_0 \notin K_f(u)$ we will show that for a smooth function φ with support on a neighborhood of x_0 exists, such that $\mathcal{F}(f^*u\varphi)$ is rapidly decreasing on W^c (which is a neighborhood of ξ_0).

We choose φ to be a smooth function with $\text{supp}(\varphi) \subseteq X'$. For all $\varepsilon < \varepsilon'$ the identity $(f_\varepsilon^*)u_\varepsilon \varphi = f_\varepsilon^*(\chi u_\varepsilon) \varphi$ holds, since the functions χ and φ where choosen such that $(\chi \circ f_\varepsilon) \cdot \varphi \equiv 1$ for $\varepsilon < \varepsilon_1$.

Set $\tilde{V} := \{\eta \in S^{m-1} \mid \exists \eta_1 \in V, \lambda \in \mathbb{R}_+ : \eta = \lambda \cdot \eta_1\}$ and $\tilde{W} := \{\xi \in S^{m-1} \mid \exists \xi \in W, \lambda \in \mathbb{R}_+ : \xi = \lambda \cdot \xi_1\}$. Obviously $\mathcal{F}(u\chi)$ is rapidly decreasing on V^c . We have that

$$\begin{aligned} |\mathcal{F}((f^*u)_\varepsilon \varphi)(\xi)| &= |\mathcal{F}((f^*\chi u)_\varepsilon \varphi)(\xi)| \\ &= \left| \int_{\mathbb{R}^m} \widehat{\chi u_\varepsilon}(\eta) \left(\int_{\mathbb{R}^n} \exp(i\langle f_\varepsilon(x), \eta \rangle - i\langle x, \xi \rangle) \varphi(x) dx \right) d\eta \right| \\ &= \left| \int_{\mathbb{R}^m} \widehat{\chi u_\varepsilon}(\eta) I_\varepsilon(\xi, \eta) d\eta \right| = \left| \int_{\tilde{V}} \widehat{\chi u_\varepsilon}(\eta) I_\varepsilon(\xi, \eta) d\eta \right| + \left| \int_{\tilde{V}^c} \widehat{\chi u_\varepsilon}(\eta) I_\varepsilon(\xi, \eta) d\eta \right|, \end{aligned}$$

where we have set

$$I_\varepsilon(\xi, \eta) := \int \exp(i\langle f_\varepsilon(x), \eta \rangle - i\langle x, \xi \rangle) \varphi(x) dx.$$

We intend to apply the stationary phase theorem 6.3 (cf. the Appendix) with

$$\begin{aligned} \omega &:= |\xi| + |\eta| \\ \phi_\varepsilon &:= \langle f_\varepsilon(x), \frac{\eta}{|\eta| + |\xi|} \rangle - \langle x, \frac{\xi}{|\eta| + |\xi|} \rangle. \end{aligned}$$

Thus we have to find a bound for the gradient of the phase function $|\phi'_\varepsilon(x)| = |{}^T df_\varepsilon(x) \alpha \eta_1 - (1 - \alpha) \xi_1|$ from below for all $\alpha \in [0, 1]$, $\eta_1 = \eta/|\eta|$, $\xi_1 = \xi/|\xi|$ and $x \in X$. By optimization in the parameter α and using the notation of Lemma 3.7 we can bound the expression

$$\inf_{x \in X, \eta_1 \in V, \xi_1 \in W, \alpha \in [0, 1]} |{}^T df_\varepsilon(x) \alpha \eta_1 - (1 - \alpha) \xi_1|$$

from below by

$$\begin{aligned} &\inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} \frac{|{}^T df_\varepsilon(x) \eta_1|}{|{}^T df_\varepsilon(x) \eta_1 + \xi_1|} \sqrt{\left| 1 - \left\langle M_\varepsilon(x, \eta_1), \xi_1 \right\rangle^2 \right|} \\ &= \inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} \frac{1}{2} \frac{|{}^T df_\varepsilon(x) \eta_1|}{|{}^T df_\varepsilon(x) \eta_1 + \xi_1|} |M_\varepsilon(x, \eta_1) - \xi_1| \\ &\geq \inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} \frac{1}{4} \frac{|{}^T df_\varepsilon(x) \eta_1|}{|{}^T df_\varepsilon(x) \eta_1| + 1} \inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} |M_\varepsilon(x, \eta_1) - \xi_1| \\ &\geq C \sigma_\varepsilon^{-1} \inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} |M_\varepsilon(x, \eta_1) - \xi_1|, \end{aligned}$$

where σ_ε is the slow-scaling net from the Definition 3.4. According to Lemma 4.3 we have that

$$d_\varepsilon := \inf_{(x, \eta_1, \xi_1) \in X \times V \times W^c} |M_\varepsilon(x, \eta_1) - \xi_1| > d > 0$$

for $\varepsilon < \varepsilon'$, where d is a positive constant. Thus the gradient of the phase function is uniformly bounded from below by $|\phi'_\varepsilon(x)| \geq Cd \cdot \sigma_\varepsilon$ for all $(x, \xi, \eta) \in X \times \tilde{V} \times \tilde{W}^c$. The stationary phase theorem 6.3 yields

$$|I_\varepsilon(\xi, \eta)| \leq C_q \varepsilon^{-1} (1 + |\xi| + |\eta|)^{-q} \quad (5)$$

for all $q \in \mathbb{N}_0$. Note that we use C, C_p, C_q and C_k, l as generic constants. In the case where $(x, \eta, \xi) \in X \times \tilde{V}^c \times \tilde{W}^c$ the stationary phase theorem (now with the phase function $\exp(-i\langle x, \xi \rangle)$) gives

$$\begin{aligned} |\xi|^k |I_\varepsilon(\xi, \eta)| &= \left| \int \exp(-i\langle x, \xi \rangle) \exp(i\langle f_\varepsilon(x), \eta \rangle) \varphi(x) dx \right| \\ &\sum_{|\alpha| \leq k} |D_x^\alpha (\exp(i|\eta| \langle f_\varepsilon(x), \eta/|\eta| \rangle) \varphi)| \end{aligned} \quad (6)$$

and by repeated use of the chain rule we obtain the estimate

$$\begin{aligned} &\leq \sum_{|\alpha| \leq k} \sum_{\beta \leq \alpha} c_1(\beta) \sup_{x \in X_0} |\partial^{\alpha-\beta} \varphi(x)| \sum_{l=1}^{|\beta|} |\eta|^l \cdot \\ &\sum_{\gamma_1 + \dots + \gamma_l = \beta} d(\gamma_1, \dots, \gamma_l) \prod_{1 \leq i \leq l} \sup_{x \in X_0} |\langle \partial^{\gamma_i} f_\varepsilon(x), \eta/|\eta| \rangle| \end{aligned}$$

where $\gamma_1 + \gamma_2 + \dots + \gamma_l$ denotes a partition of the multiindex β in exactly l multiindices, that add up componentwise to β . Using the notation $|g|_k := \max_{|\alpha|=k} |\partial^\alpha g(x)|$, we can bound the expression by

$$\begin{aligned} &\leq C_{k,2} \sum_{|\alpha| \leq k} \sum_{\beta \leq \alpha} \sum_{l=0}^{|\beta|} |\eta|^l \max_{\gamma_1 + \dots + \gamma_l = \beta} \left(\prod_{1 \leq i \leq l} \sup_{x \in X, \eta \in \tilde{V}^c} |\langle f_\varepsilon(x), \eta/|\eta| \rangle|_{|\gamma_i|} \right) \\ &\leq C_{k,2} \sum_{|\alpha| \leq k} \sum_{\beta \leq \alpha} \sum_{l=0}^{|\beta|} |\eta|^l \max_{0 \leq j \leq |\beta| - l + 1} \left(\sup_{x \in X, \eta \in \tilde{V}^c} |\langle f_\varepsilon(x), \eta/|\eta| \rangle|_j \right)^l \\ &\leq C_{k,3} (1 + |\eta|)^k \max_{l=1}^k \max_{0 \leq j \leq k-l+1} \left(\sup_{x \in X, \eta \in \tilde{V}^c} |\langle f_\varepsilon(x), \eta/|\eta| \rangle|_j \right)^l. \end{aligned}$$

Since $f_\varepsilon(x)$ is of slow-scale in all derivatives at x_0 , it holds that for all $j \in \mathbb{N}_0$ there exists constants C_j and slow-scaled nets $r_{\varepsilon,j}$ such that

$$\sup_{(x, \eta_1) \in X \times V^c} |\langle f_\varepsilon(x), \eta_1 \rangle|_j \leq C_j r_{\varepsilon,j}$$

holds for small ε . Summing up the estimates of the form (6) give that there exists constants C_p such that

$$|I_\varepsilon(\xi, \eta)| \leq C_p (1 + |\eta|)^{-p} \varepsilon^{-1} (1 + |\xi|)^p \quad (7)$$

holds for all $p \in \mathbb{N}_0$ and $(\eta, \xi) \in \widetilde{V}^c \times \widetilde{W}$. We observe that

$$\begin{aligned} |\mathcal{F}((f^*u)_\varepsilon\varphi)(\xi)| &= \left| \int_{\mathbb{R}^m} \widehat{\chi u_\varepsilon}(\eta) I_\varepsilon(\xi, \eta) d\eta \right| \\ &\leq C_q \varepsilon^{-1} \int_{\widetilde{V}} |\widehat{\chi u_\varepsilon}(\eta)| (1 + |\xi| + |\eta|)^{-q} d\eta \\ &\quad + C_p (1 + |\xi|)^{-p} \varepsilon^{-1} \int_{\widetilde{V}^c} |\widehat{\chi u_\varepsilon}(\eta)| (1 + |\eta|)^p d\eta \end{aligned}$$

holds and using (5) and (7) leads to the upper bound

$$\begin{aligned} &C_q \varepsilon^{-1} \sup_{\eta \in \widetilde{V}} |(1 + |\eta|)^{-k} \widehat{\chi u_\varepsilon}(\eta)| \int_{\widetilde{V}} (1 + |\xi| + |\eta|)^{k-q} d\eta \\ &+ C_p (1 + |\xi|)^{-p} \varepsilon^{-1-n} \int_{\widetilde{V}^c} (1 + |\eta|)^{p-l} d\eta. \end{aligned}$$

Finally we set $k := q - p - m$ and $l := p + n - 1$ and obtain

$$\begin{aligned} &C_q \varepsilon^{-1} \sup_{\eta \in \widetilde{V}} |(1 + |\eta|)^{-k} \widehat{\chi u_\varepsilon}(\eta)| \int_{\widetilde{V}} (1 + |\xi| + |\eta|)^{k-q} d\eta \\ &+ C_p (1 + |\xi|)^{-p} \varepsilon^{-1-n} \int_{\widetilde{V}^c} (1 + |\eta|)^{p-l} d\eta \\ &\leq C_{p,q,m,n} \varepsilon^{-1-n} (1 + |\xi|)^{-p} \end{aligned}$$

for all $\xi \in \widetilde{W}^c$ and $C_{p,q,m,n}$ some constant depending on p, q, m and n . It follows that $(x_0, \xi_0) \notin \text{WF}(f^*u)$. $\square$

5 Examples

Example 5.1 (Multiplication of Colombeau functions). *This example was presented in [10, Example 4.2] in order to show that an inclusion relation like in [7, Theorem 8.2.10] for the wave front set of a product of distributions, cannot be extended to Colombeau function with wavefront sets in unfavorable position.*

Consider the Colombeau functions u and v defined by

$$\begin{aligned} u_\varepsilon &:= \varepsilon^{-1} \rho(\varepsilon^{-1}(x + \gamma_\varepsilon y)) \\ v_\varepsilon &:= \varepsilon^{-1} \rho(\varepsilon^{-1}(x - \gamma_\varepsilon y)), \end{aligned}$$

where γ_ε is some net with $\lim_{\varepsilon \rightarrow 0} \gamma_\varepsilon = 0$. Note that in [10, Example 4.2] $\gamma_\varepsilon := \varepsilon^{1/2}$. These Colombeau functions are both associated to $\delta(x) \otimes 1(y)$ and the wavefront sets $\text{WF}(u) = \text{WF}(v) = \{0\} \times \mathbb{R} \times \{(\pm 1, 0)\}$ are in an unfavorable position.

We are going to apply Theorem (4.4) in order to calculate $\text{WF}(u \cdot v)$. First we rewrite $u \cdot v = f^* \iota(\delta)$, where $f_\varepsilon(x, y) = (x + \gamma_\varepsilon y, x - \gamma_\varepsilon y)$ with γ_ε some net tending

to zero. In Example (3.5) we already showed that $D_f = \mathbb{R}^2 \times S^1 \setminus (\pm 1, \mp 1)$. From [9, Theorem 15] it follows that $WF(\iota(\delta)) = \{(0, 0)\} \times S^1$. The wavefront unfavorable support of f with respect to $WF(\iota(\delta))$ according to Definition 3.4 is

$$U_f((0, 0) \times S^1) = \{(0, 0)\}.$$

Since $f_\varepsilon(x, y)$ is of slow-scale in all derivatives at all $x \in \mathbb{R}^2$, it follows that $S_f^c = \emptyset$. Now Theorem 4.4 gives that

$$WF(u \cdot v) \subseteq (0, 0) \times S^1,$$

which is consistent with the result in [10, Example 4.2].

Example 5.2 (Hurd-Sattinger). *Let us consider the initial value problem*

$$\begin{aligned} \partial_t u + \Theta \partial_x u + \Theta' u &= 0 \\ u(0, x) &= u_0 \in \mathcal{G}(\mathbb{R}), \end{aligned} \tag{8}$$

where $\Theta \in \mathcal{G}(\mathbb{R}^2)$ is defined by $\Theta_\varepsilon(x) = \rho_{\gamma_\varepsilon} * H(-\cdot)$ with $\rho_{\gamma_\varepsilon} = \frac{1}{\gamma_\varepsilon} \rho(\cdot/\gamma_\varepsilon)$ where $\rho \in \mathcal{S}(\mathbb{R})$, $\rho \geq 0$ and $\int \rho(x) dx = 1$ and $\gamma_\varepsilon = \log(1/\varepsilon)$ is a net of slow-scale. For the initial value we choose $u_0 := \iota(\delta_{-s_0})$ a delta like singularity at $-s_0$ (for a positive $s_0 > 0$). The Hurd-Sattinger example was first given in [12] (it was shown that it is not solvable in L^1_{loc} , when distributional products are employed). It was further investigated in [9] with methods from Colombeau theory. In [5] the wavefront set WF_γ (with respect to the slow-scale net γ) of the Colombeau solution was calculated. For sake of simplicity we do only consider the standard generalized wave-front set WF , which is smaller since it neglects the singularities coming from the coefficient Θ .

We have

$$\Theta_\varepsilon(x) = \int_{x/\gamma_\varepsilon}^{\infty} \rho(z) dz.$$

Consider the ordinary differential equation for the characteristic curve, which passes through (t, x) at time t :

$$\begin{aligned} \partial_0 \sigma_\varepsilon(s; t, x) &= \Theta_\varepsilon(\sigma_\varepsilon(s; t, x)) \\ \sigma(t; t, x) &= x. \end{aligned}$$

The derivatives of the characteristics fulfill linear initial value problems: If we set $y_\varepsilon := \partial_t \sigma_\varepsilon(s; t, x)$ we obtain

$$\dot{y}_\varepsilon = \Theta_\varepsilon(\sigma_\varepsilon(s; t, x)) y_\varepsilon, \quad y_\varepsilon(t; t, x) = -\Theta_\varepsilon(x).$$

If we set $z_\varepsilon(s; t, x) := \partial_x \sigma_\varepsilon(s; t, x)$, we obtain

$$\dot{z}_\varepsilon = -\Theta_\varepsilon(\sigma_\varepsilon(s; t, x)) z_\varepsilon, \quad z_\varepsilon(t; t, x) = 1.$$

This yields

$$\partial_t \sigma_\varepsilon(s; t, x) = -\Theta_\varepsilon(\sigma_\varepsilon(s; t, x)) \text{ and } \partial_x \sigma_\varepsilon(s; t, x) = \frac{\Theta_\varepsilon(\sigma_\varepsilon(s; t, x))}{\Theta_\varepsilon(x)}.$$

If we set $f_\varepsilon(t, x) := (t, \sigma_\varepsilon(0; t, x))$ then the solution of the initial value problem (8) reads

$$u_\varepsilon = f_\varepsilon^*(1 \otimes u_{0,\varepsilon}) \cdot \frac{\Theta_\varepsilon(\sigma_\varepsilon(0; t, x))}{\Theta_\varepsilon(x)}.$$

We expect that the wave front set of u is generated from the first factor $f^*(1 \otimes u_0)$, since the second factor $\frac{\Theta_\varepsilon(\sigma_\varepsilon(0; t, x))}{\Theta_\varepsilon(x)}$ contains only slow-scale terms. The wavefront set corresponding to the initial value is $\Gamma := \text{WF}(1 \otimes u_0) = \mathbb{R} \times \{-s_0\} \times \{0\} \times \{\pm 1\}$ and the transposed Jacobian of the generalized pullback f is

$$df_\varepsilon(t, x)^T = \begin{pmatrix} 1 & -\Theta_\varepsilon(\sigma_\varepsilon(0; t, x)) \\ 0 & \frac{\Theta_\varepsilon(\sigma_\varepsilon(0; t, x))}{\Theta_\varepsilon(x)} \end{pmatrix}.$$

If the Jacobian acts on $\zeta := (0, \pm 1) \in S^1$ (which are the only irregular directions coming from the wavefront set of $1 \otimes u_0$), we obtain

$$\begin{aligned} M_\varepsilon(t, x, \zeta) &= \frac{df_\varepsilon(t, x)^T \zeta}{|df_\varepsilon(t, x)^T \zeta|} \\ &= \pm \left(\frac{-1}{\sqrt{1 + \Theta_\varepsilon(x)^{-2}}}, \frac{1}{\sqrt{1 + \Theta_\varepsilon(x)^2}} \right). \end{aligned}$$

Remarkably the result does only depend on the coefficient $\Theta_\varepsilon(x)$. Let $z_\varepsilon := (t_\varepsilon, x_\varepsilon) \rightarrow (t_0, x_0)$ be some convergent net, then we immediately get that

$$M_\varepsilon(z_\varepsilon, \zeta) \rightarrow \begin{cases} (-\zeta_1, \zeta_2) & \text{for } x_0 < 0 \\ (0, \zeta_2) & \text{for } x_0 > 0 \\ \left(-\frac{\alpha}{\sqrt{1+\alpha^2}} \zeta_1, \frac{1}{\sqrt{1+\alpha^2}} \zeta_2 \right), \alpha \in [0, 1] & \text{for } x_0 = 0. \end{cases}$$

In order to obtain estimates for the characteristic flow we note that $0 \leq \Theta_\varepsilon(z) \leq 1$, since the ρ was assumed to be positive. We conclude that

$$\sigma_\varepsilon(0; t, x) < \sigma_\varepsilon(0; t, y) : x < y \text{ and } \sigma_\varepsilon(0; p, x) < \sigma_\varepsilon(0; q, x) : q < p,$$

since $\partial_t \sigma_\varepsilon(0; t, x) < 0$ and $\partial_x \sigma_\varepsilon(0; t, x) > 0$. Thus it is clear that

$$x - t \leq \sigma_\varepsilon(0; t, x) \leq x \tag{9}$$

holds globally. Since Θ_ε is monotone and positive, we can improve these bounds using the integral representation of $\sigma_\varepsilon(0; t, x)$ by

$$x - t\Theta_\varepsilon(x - t) \leq \sigma_\varepsilon(0; t, x) \leq x - t\Theta_\varepsilon(x). \tag{10}$$

In order to determine C_{Φ_f} we have to calculate all clusterpoints of nets $(\sigma_{\tau(\varepsilon)}(0; t_\varepsilon, x_\varepsilon))_\varepsilon$ for arbitrary nets $(t_\varepsilon, x_\varepsilon)_\varepsilon$ tending to (t_0, x_0) and $\tau :]0, 1] \mapsto]0, 1]$ some map with $\tau(\varepsilon) < \varepsilon$. Assume $t_0 > 0$, then we have to distinguish three cases: If $x_0 < 0$ then we can use the estimate (10) to obtain $\lim_{\varepsilon \rightarrow 0} \sigma_{\tau(\varepsilon)}(0; t_\varepsilon, x_\varepsilon) = x_0 - t_0$. In the case $x_0 > 0$ we first solve $\sigma_{\tau(\varepsilon)}(0; t, x) = x_0/2$ in the time variable and obtain globally defined (since $\sigma_{\tau(\varepsilon)}(0; t, x)$ is monotone) family of functions $T_\varepsilon(x)$ with the property that $\sigma_{\tau(\varepsilon)}(0; T_\varepsilon(x), x) = x_0/2$. If $x > x_0/2$ we have that $T_\varepsilon(x) > 0$. Without loss of generality we assume that $x_\varepsilon > x_0/2$. Using (10) we derive the estimate

$$T_\varepsilon(x) \geq \frac{x - x_0/2}{\Theta_{\tau(\varepsilon)}(x_0/2)}.$$

It yields that $\lim_{\varepsilon \rightarrow 0} T_\varepsilon(x_\varepsilon) = +\infty$, so $t_\varepsilon < T_\varepsilon(x_\varepsilon)$ holds for small ε . We obtain

$$x_\varepsilon - t_\varepsilon \Theta_\varepsilon(x_0/2) \leq \sigma_\varepsilon(0; t_\varepsilon, x_\varepsilon) \leq x_\varepsilon - t_\varepsilon \Theta_\varepsilon(x_\varepsilon)$$

as ε gets small. It follows that $\lim_{\varepsilon \rightarrow 0} \sigma_{\tau(\varepsilon)}(0; t_\varepsilon, x_\varepsilon) = x_0$.

The case $x_0 = 0$ is a bit more difficult: We proceed with an idea from the proof of [13, Proposition 3]. First off all we can observe that $-t \leq \lim_{\varepsilon \rightarrow 0} \sigma_{\tau(\varepsilon)}(0; t, x) \leq 0$. Let $(a_\varepsilon)_\varepsilon$ be a strictly positive net with $\lim_{\varepsilon \rightarrow 0} a_\varepsilon = 0$ and $\lim_{\varepsilon \rightarrow 0} \Theta_{\tau(\varepsilon)}(a_\varepsilon)$. Furthermore we define $t_\varepsilon := t_0 + a_\varepsilon$. In the next step we construct nets $(x_{\varepsilon, \alpha})_\varepsilon$ for each $\alpha \in [-t_0, 0]$ that converge to zero and have the property that $\lim_{\varepsilon \rightarrow 0} \sigma_{\tau(\varepsilon)}(0; t_\varepsilon, x_{\varepsilon, \alpha}) = \alpha$. We set $x_{\varepsilon, \alpha} := \sigma_{\tau(\varepsilon)}(t_\varepsilon; 0, \alpha)$. Let $S_\varepsilon(\alpha)$ be the function, that solves $\sigma_{\tau(\varepsilon)}(S_\varepsilon(\alpha); 0, \alpha) = a_\varepsilon$ globally. It exists because $\partial_x \sigma_\varepsilon(t; 0, x)$ is strictly positive and $\sigma_\varepsilon(t; 0, x)$ is monotone. Since $a_\varepsilon > 0$ and $\alpha \in [-t_0, 0]$ we have that $S_\varepsilon(\alpha) \geq 0$. Furthermore we can apply (9) to obtain $S_\varepsilon(\alpha) \leq -\alpha + a_\varepsilon$. For all $t \geq S_\varepsilon(\alpha)$ it follows that

$$a_\varepsilon \leq \sigma_{\tau(\varepsilon)}(t; 0, \alpha) \leq a_\varepsilon + (t - S_\varepsilon(\alpha))\Theta_{\tau(\varepsilon)}(a_\varepsilon) \quad (11)$$


and we can conclude that $\lim_{\varepsilon \rightarrow 0} \sigma_{\tau(\varepsilon)}(t_\varepsilon; 0, \alpha) = 0$, since a_ε was choosen such that $\lim_{\varepsilon \rightarrow 0} \Theta_{\tau(\varepsilon)}(a_\varepsilon) = 0$ and $t_\varepsilon := t_0 + a_\varepsilon \geq -\alpha + a_\varepsilon \geq S_\varepsilon(\alpha)$ for ε small enough. Finally we have that the generalized graph of the phasefunction ϕ_f coming from the characteristic flow $f_\varepsilon(t, x) = (t, \sigma_\varepsilon(0; t, x))$ is

$$\begin{aligned} C_{\phi_f} \cap \tilde{\Gamma} &= \{(t, t - s_0, 0, \pm 1; 0, -s_0, \mp 1, \pm 1) \mid t < s_0\} \cup \\ &\quad \{(s_0, 0, 0, \pm 1; 0, -s_0, \mp \frac{\alpha}{\sqrt{1+\alpha^2}}, \pm \frac{1}{\sqrt{1+\alpha^2}}) \mid \alpha \in [0, 1]\} \cup \\ &\quad \{(t, 0, 0, \pm 1; t, -s_0, 0, \pm 1) \mid t > s_0, \alpha \in [0, 1]\} \end{aligned}$$

where $\tilde{\Gamma} := \{(x, \eta, y, \xi) \in \mathbb{R} \times S^1 \times \mathbb{R} \times S^1 \mid (y, \eta) \in \Gamma\}$. By Theorem 4.4 we obtain

$$\begin{aligned} WF(u) = WF(f^*(1 \otimes u_0)) &\subseteq \pi_2(C_{\phi_f} \cap \tilde{\Gamma}) = \\ &\quad \{(t, t - s_0; \mp 1, \pm 1) \mid t < s_0\} \cup \\ &\quad \{(s_0, 0; \mp \frac{\alpha}{\sqrt{1+\alpha^2}}, \pm \frac{1}{\sqrt{1+\alpha^2}}) \mid \alpha \in [0, 1]\} \cup \\ &\quad \{(t, 0; 0, \pm 1) \mid t > s_0, \alpha \in [0, 1]\}. \end{aligned}$$

Figure 1: Estimated wave front set of the generalized solution of the Hurd-Sattinger partial differential equation.



6 Appendix: Stationary Phase theorem

Definition 6.1. Suppose that $g \in \mathcal{G}(\Omega)$ and let $(g_\varepsilon)_\varepsilon$ be some representative, then we introduce the notation

$$|g_\varepsilon|_k := \sum_{|\alpha|=k} |\partial^\alpha g_\varepsilon|$$

and

$$\begin{aligned} \mu_{K,k,\varepsilon}(g) &:= \sup_K |g_\varepsilon|_k \\ \mu_{K,k,\varepsilon}^*(g) &:= \max_{l \leq k} \mu_{K,l,\varepsilon}(g). \end{aligned}$$

Lemma 6.2. Suppose that $g \in \mathcal{G}(\Omega)$ with an non-negative representative $(g_\varepsilon)_\varepsilon$, such that there exists $\varepsilon' > 0$, with $g_\varepsilon(x) \geq 0$ for $\varepsilon < \varepsilon'$ and all $x \in \Omega$. Then it follows that for any $K \subset\subset \Omega$ some compact set there exist some C, N such that

$$\delta^2 \sum_{|\alpha|=1} |\partial^\alpha g_\varepsilon(x)| \leq C \sqrt{g_\varepsilon(x)} \sqrt{\mu_{M,2,\varepsilon}^*(g)}$$

holds for all $x \in K$ and M is some compact set with $K \subset\subset M^\circ$.

Proof. Let K be some compact set and $M \subset\subset \Omega$ such that $K \subset\subset M^\circ$. We use the notation $B_\delta(K) := \{x \in \Omega \mid \exists x_0 \in K : |x - x_0| \leq \delta\}$. Then we can find some $\delta > 0$ such that $B_\delta(K) \subseteq M$ for some $\delta > 0$. Then we have for all $x \in K$

that $B_\delta(x_0) \subseteq M$. Choose some $x_0 \in K$. By Taylor's formula we obtain for any $x = x_0 + \alpha v$ with $\alpha \in [0, \delta]$ and $|v| = 1$ (note that $x_0 + \alpha v \in B_\delta(x_0) \subseteq M$), that

$$0 \leq g_\varepsilon(x) \leq g_\varepsilon(x_0) + \partial_v g_\varepsilon(x_0)\alpha + \frac{1}{2}m_\varepsilon\alpha^2,$$

where

$$m_\varepsilon(v) := \max(\sup_M |\partial_v^2 g_\varepsilon|, 2g_\varepsilon(x_0)\delta^{-2}).$$

So we can obtain the estimate

$$|\partial_v g_\varepsilon(x_0)| \leq \alpha^{-1}g_\varepsilon(x_0) + \frac{\alpha}{2}m_\varepsilon(v).$$

Now we distinguish two cases: Let $\delta^2 m_\varepsilon \leq 4g_\varepsilon(x_0)$, then we set $\alpha = \delta$ and obtain

$$\begin{aligned} \delta |\partial_v g_\varepsilon(x_0)| &\leq g_\varepsilon(x_0) + \frac{\delta^2}{2}m_\varepsilon(v) \\ &= \sqrt{(g_\varepsilon(x_0) + \frac{\delta^2}{2}m_\varepsilon(v))^2} \\ &= \sqrt{(g_\varepsilon(x_0)^2 + \delta^2 g_\varepsilon(x_0)m_\varepsilon(v) + \frac{\delta^4}{4}m_\varepsilon(v)^2)} \\ &\leq \sqrt{(g_\varepsilon(x_0)^2 + \delta^2 g_\varepsilon(x_0)m_\varepsilon(v) + \delta^2 m_\varepsilon(v)g_\varepsilon(x_0))} \\ &= \sqrt{g_\varepsilon(x_0)}\sqrt{g_\varepsilon(x_0) + 2\delta^2 m_\varepsilon(v)}. \end{aligned}$$

In the case where $\delta^2 m_\varepsilon(v) > 4g_\varepsilon(x_0)$ we set $\alpha := \sqrt{2g_\varepsilon(x_0)/m_\varepsilon(v)} < \delta$ and obtain

$$\begin{aligned} |\partial_v g_\varepsilon(x_0)| &\leq \sqrt{m_\varepsilon/2}\sqrt{g_\varepsilon(x_0)} + \frac{\sqrt{2g_\varepsilon(x_0)m_\varepsilon(v)}}{2\sqrt{m_\varepsilon(v)}} \\ &= \sqrt{2m_\varepsilon(v)}\sqrt{g_\varepsilon(x_0)} \end{aligned}$$

We can finally conclude that the estimate

$$\delta^2 |\partial_v g_\varepsilon(x_0)|^2 \leq g_\varepsilon(x_0)(g_\varepsilon(x_0) + 2\delta^2 m_\varepsilon(v)) \leq g_\varepsilon(x_0)(\sup_{x \in K} g_\varepsilon(x) + 2\delta^2 m_\varepsilon(v))$$

holds for all $x_0 \in K$, where $m_\varepsilon(v)$ is independant of x_0 . $\square$

Theorem 6.3 (Stationary phase theorem). *Let $u \in \mathcal{G}_c(\Omega)$ with support $K \subset\subset \Omega$ and $\phi_\varepsilon \in \mathcal{E}(\Omega)$ with the property that there exists an $\varepsilon_0 > 0$ and $m \in \mathbb{N}$ such that*

$$\inf_{x \in K} (|\phi'_\varepsilon(x)|) \geq \lambda_\varepsilon \quad \text{for all } \varepsilon \leq \varepsilon_0 \quad (12)$$

holds, with λ_ε some net tending to zero. Then we have that

$$(v_\varepsilon)_\varepsilon := \left(\int u_\varepsilon(x) \exp(i\omega\phi_\varepsilon(x)) dx \right)_\varepsilon + \mathcal{N}(\Omega) \quad (13)$$

is a Colombeau function in the ω variable and it is bounded by

$$\omega^k |v_\varepsilon(\omega)| \leq L_{k,\varepsilon} \lambda_\varepsilon^{-k} \sum_{|\alpha| \leq k} \sup_K |D^\alpha u_\varepsilon|, \quad (14)$$

where

$$L_{k,\varepsilon} := C_k \max \{1, \mu_{M,k,\varepsilon}^*(\phi_\varepsilon)^{2k^2}\}$$

and M is some compact set with $K \subset M^\circ$.

Proof. It is obvious that (13) is a well-defined Colombeau function. The proof follows closely the proof of classical stationary phase theorem in [7, Theorem 7.7.1]. By $N_\varepsilon(x) := |\phi'_\varepsilon(x)|^2$ we denote the square of the norm of the gradient of the phase function. Let

$$u_{\nu,\varepsilon} := N_\varepsilon^{-1} \frac{\partial \phi_\varepsilon}{\partial x_\nu} u_\varepsilon$$

and since

$$i\omega \frac{\partial \phi_\varepsilon}{\partial x_\nu} \exp(i\omega \phi_\varepsilon) = \partial_\nu \exp(i\omega \phi_\varepsilon)$$

we obtain after an integration by parts

$$\int u_\varepsilon \exp(i\omega \phi_\varepsilon) dx = \frac{i}{\omega} \sum_\nu \int (\partial_\nu u_{\nu,\varepsilon}) \exp(i\omega \phi_\varepsilon) dx$$

(using the notation introduced in Definition 6.1). We prove by induction: For $k = 0$ we have the obvious bound

$$\left| \int u_\varepsilon \exp(i\omega \phi_\varepsilon) dx \right| \leq C \sup_K |u_\varepsilon(x)|.$$

Assume that the bound (14) holds for power $k - 1$, then we have that

$$\begin{aligned} \omega^k \left| \int u_\varepsilon \exp(i\omega \phi_\varepsilon) dx \right| &= \omega^{k-1} \left| \sum_\nu \int (\partial_\nu u_{\nu,\varepsilon}) \exp(i\omega \phi_\varepsilon) dx \right| \\ &\leq L_{k-1,\varepsilon} \sum_{m=0}^{k-1} \sup_K \left(\sum_{\nu=1}^n |u_{\nu,\varepsilon}|_{\mu+1} N_\varepsilon^{m/2-k+1} \right) \end{aligned}$$

holds. In the next step we are going to show that

$$N_\varepsilon^{\frac{1}{2}} \sum_\nu |u_{\nu,\varepsilon}|_m \leq M_{m,\varepsilon} \sum_{r=0}^m |u_\varepsilon|_r N_\varepsilon^{\frac{r-m}{2}} \quad (15)$$

holds. Again we are proving by induction: For $m = 0$ we have

$$N_\varepsilon^{\frac{1}{2}} \sum_\nu |u_{\nu,\varepsilon}| = |u_\varepsilon| \sum_\nu N_\varepsilon^{-\frac{1}{2}} \left| \frac{\partial \phi_\varepsilon}{\partial x_\nu} \right| \leq n |u_\varepsilon| \quad (16)$$

and let us now assume that (15) holds up to $m - 1$. Let α be any multi-index with $|\alpha| = m$. We apply ∂^α on

$$N_\varepsilon u_{\nu,\varepsilon} = u_\varepsilon \frac{\partial \phi_\varepsilon}{\partial x_\nu}$$

and obtain

$$\partial^\alpha (N_\varepsilon u_{\nu,\varepsilon}) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (\partial^\beta u_\varepsilon) (\partial^{\alpha-\beta+e_\nu} \phi_\varepsilon).$$

It follows that

$$\begin{aligned} |N_\varepsilon \partial^\alpha u_\nu^\varepsilon| &= \left| - \sum_{\beta \neq 0, \beta \leq \alpha} \binom{\alpha}{\beta} \partial^\beta N_\varepsilon \partial^{\alpha-\beta} u_{\nu,\varepsilon} + \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \partial^\beta u_\varepsilon \partial^{\alpha-\beta+e_\nu} \phi_\varepsilon \right| \\ &= C \left(\sum_{l=1}^m |N_\varepsilon|_l |u_{\nu,\varepsilon}|_{m-l} + \sum_{l=0}^m |\phi_\varepsilon|_{m-l+1} |u_\varepsilon|_l \right) \\ &= C (|N_\varepsilon|_1 |u_{\nu,\varepsilon}|_{m-1} + |\phi_\varepsilon|_1 |u_\varepsilon|_m \\ &\quad + \max_{l=0}^{m-2} (\sup |N_\varepsilon|_{m-l}, \sup |\phi_\varepsilon|_{m-l+1}) \sum_{l=0}^{m-2} (|u_{\nu,\varepsilon}|_l + |u_\varepsilon|_l) \\ &\quad + |u_\varepsilon|_{m-1} |\phi_\varepsilon|_2). \end{aligned} \tag{17}$$

Now we can apply Lemma 6.2 to bound $|N_\varepsilon|_1$. This yields

$$\begin{aligned} |N_\varepsilon(x)|_1 &= \sum_{|\alpha|=1} |\partial^\alpha N_\varepsilon(x)| \leq C_1 \sqrt{N_\varepsilon(x)} \sqrt{\mu_{L,2,\varepsilon}^*(N_\varepsilon)} \\ |\phi_\varepsilon|_1 &\leq C_2 \sqrt{N_\varepsilon(x)} \end{aligned}$$

and we can verify that

$$\begin{aligned} |\partial^\alpha N_\varepsilon| &= \left| \sum_{i=1}^n \partial^\alpha (\partial^{e_i} \phi_\varepsilon(x))^2 \right| \\ &\leq K_1 \sum_{i=1}^n \sum_{\beta \leq \alpha} |\partial^{\beta+e_i} \phi_\varepsilon(x)| |\partial^{\alpha-\beta+e_i} \phi_\varepsilon(x)| \\ &\leq K_2 \sum_{k \leq |\alpha|} |\phi_\varepsilon(x)|_{|\alpha|-k+1} |\phi_\varepsilon(x)|_{k+1} \end{aligned}$$

holds. This yields

$$\begin{aligned} \sup_K |N_\varepsilon|_l &\leq K_3 \sum_{k \leq l} \sup_K |\phi_\varepsilon(x)|_{l-k+1} \sup_K |\phi_\varepsilon(x)|_{k+1} \\ &\leq K_4 \mu_{K,l+1,\varepsilon}^*(\phi_\varepsilon)^2. \end{aligned}$$

We introduce

$$\begin{aligned}\sigma_{\varepsilon, m} &:= \max_{l=2}^m (\sup |N_\varepsilon|_l, \sup |\phi_\varepsilon|_{l+1}) \\ &\leq \max (K_4 \mu_{K, m+1, \varepsilon}^* (\phi_\varepsilon)^2, \mu_{K, m+1, \varepsilon}^* (\phi_\varepsilon))\end{aligned}$$

in order to simplify the notation. Using these bounds we can estimate (17) by

$$\begin{aligned}& C \left(|N_\varepsilon|^{1/2} C_1 \sqrt{\mu_{L, 2, \varepsilon}^* (N_\varepsilon)} |u_{\nu, \varepsilon}|_{m-1} + C_2 |N_\varepsilon|^{1/2} |u_\varepsilon|_m \right. \\ & \quad \left. + C_3 \sigma_\varepsilon^{(m-2)} \left(\sum_{l=1}^{m-2} (|u_{m, \varepsilon}|_l + |u_\varepsilon|_l) \right) + |\phi_\varepsilon|_2 |u_\varepsilon|_{m-1} \right) \\ & \leq C_5 \sigma_\varepsilon^{(m-2)} \left(|N_\varepsilon|^{1/2} |u_{\nu, \varepsilon}|_{m-1} + |N_\varepsilon|^{1/2} |u_\varepsilon|_m + \sum_{l=1}^{m-2} |u_{\nu, \varepsilon}|_l + \sum_{l=1}^{m-1} |u_\varepsilon|_{m-1} \right)\end{aligned}$$

and the induction hypothesis (15) for $m-1$ gives

$$\begin{aligned}& \leq C_5 \sigma_\varepsilon^{(m)}. \\ & \left(M_{m-1, \varepsilon} \sum_{r=0}^{m-1} |u_\varepsilon|_r N_\varepsilon^{\frac{r-m+1}{2}} + |N_\varepsilon|^{1/2} |u_\varepsilon|_m \right. \\ & \quad \left. + \sum_{l=1}^{m-2} M_{l, \varepsilon} \sum_{r=0}^l |u_\varepsilon|_r N_\varepsilon^{\frac{r-l-1}{2}} + \sum_{l=1}^{m-1} |u_\varepsilon|_l \right) \\ & \leq C_5 \sigma_\varepsilon^{(m)}. \\ & \left(M_{m-1, \varepsilon} \sum_{r=0}^m |u_\varepsilon|_r N_\varepsilon^{\frac{r-m+1}{2}} + \sum_{l=0}^{m-2} \max \{1, M_{l, \varepsilon}\} \sum_{r=0}^{l+1} |u_\varepsilon|_r N_\varepsilon^{\frac{r-l-1}{2}} \right) \\ & \leq C_5 \sigma_\varepsilon^{(m)} \left(\sum_{l=0}^{m-1} \max \{1, M_{l, \varepsilon}\} \right) \sum_{r=0}^{l+1} |u_\varepsilon|_r N_\varepsilon^{\frac{r-l-1}{2}} \\ & \leq C_6 \sigma_\varepsilon^{(m)} \max_{l=0}^{m-1} \max \{1, M_{l, \varepsilon}\} \sum_{r=0}^m |u_\varepsilon|_r N_\varepsilon^{\frac{r-m-1}{2}}.\end{aligned}$$

It follows immediately that

$$\sum_{\nu} |u_{\nu, \varepsilon}|_m \leq M_{m, \varepsilon} \sum_{r=0}^m |u|_r N_\varepsilon^{\frac{r-m}{2}} \quad (18)$$

holds, where $M_{m, \varepsilon}$ is defined by

$$M_{m, \varepsilon} := \sigma_\varepsilon^{(m)} \max_{l=0}^{m-1} \max \{1, M_{l, \varepsilon}\} = C \sigma_\varepsilon^{(m)} \Pi_{i=1}^{m-1} \max \{\sigma_\varepsilon^{(i)}, 1\}$$

is a generalized number. We are finally able to estimate (6) from above by

$$\begin{aligned} & L_{k-1,\varepsilon} \sum_{m=0}^{k-1} \sup_K \left(M_{m+1,\varepsilon} \sum_{r=0}^{m+1} |u_\varepsilon|_r N_\varepsilon^{\frac{r-m-1}{2}} N_\varepsilon^{\frac{m+1}{2}-k} \right) \\ & \leq L_{k-1,\varepsilon} M_{k,\varepsilon} (k-1) \sum_{r=0}^k \sup_K \left(|u_\varepsilon|_r N_\varepsilon^{\frac{r}{2}-k} \right). \end{aligned}$$

The generalized constant $L_{k,\varepsilon}$ is recursively defined by

$$\begin{aligned} L_{k,\varepsilon} &:= L_{k-1,\varepsilon} M_{k,\varepsilon} (k-1) = (k-1)! \prod_{l=1}^k M_{l,\varepsilon} \\ &= C(k-1)! \prod_{l=1}^k \sigma_\varepsilon^{(l)} \prod_{i=1}^{l-1} \max\{\sigma_\varepsilon^{(i)}, 1\} \\ &\leq C_k \left(\prod_{l=1}^k \max\{1, \mu_{L,l,\varepsilon}^*(\phi_\varepsilon)^2\} \right) \prod_{l=1}^k \prod_{i=1}^{l-1} \max\{1, \mu_{L,i,\varepsilon}^*(\phi_\varepsilon)^2\} \\ &\leq C_k \max\{1, \mu_{L,k,\varepsilon}^*(\phi_\varepsilon)^{2k^2}\} \end{aligned}$$

Note that for $0 \leq r \leq k$ we have that

$$N_\varepsilon^{\frac{r}{2}-k} \leq \min\{1, \inf_K |\phi'_\varepsilon(x)|^{-k}\} \leq \lambda_\varepsilon^{-k}$$

holds. □

References

- [1] J. F. Colombeau. *New generalized functions and multiplication of distributions*. North-Holland, Amsterdam, 1984.
- [2] J. F. Colombeau. *Elementary introduction to new generalized functions*. North-Holland, 1985.
- [3] N. Dapić, S. Pilipović, and D. Scarpalézos. *The Linear Theory of Colombeau Generalized Functions*. Pitman Research Notes in Mathematics, Vol. 385, Longmann, Harlow, 1998.
- [4] N. Dapić, S. Pilipović, and D. Scarpalézos. Microlocal analysis of Colombeau's generalized functions: propagation of singularities. *Jour. d'Analyse Math.*, 75:51–66, 1998.
- [5] C. Garetto and G. Hörmann. On duality theory and pseudodifferential techniques for colombeau algebras: generalized delta functionals, kernels and wave front sets. *Jour. Serbian Acad. Sci.*, 2005. to appear.

- [6] M. Grosser, M. Kunzinger, M. Oberguggenberger, and R. Steinbauer. *Geometric theory of generalized functions*. Kluwer, Dordrecht, 2001.
- [7] L. Hörmander. *The analysis of linear partial differential operators*, volume I. Springer-Verlag, second edition, 1990.
- [8] G. Hörmann. Integration and microlocal analysis in Colombeau algebras. *J. Math. Anal. Appl.*, 239:332–348, 1999.
- [9] G. Hörmann and M. V. de Hoop. Microlocal analysis and global solutions of some hyperbolic equations with discontinuous coefficients. *Acta Appl. Math.*, 67:173–224, 2001.
- [10] G. Hörmann and M. Kunzinger. Microlocal analysis of basic operations in Colombeau algebras. *J. Math. Anal. Appl.*, 261:254–270, 2001.
- [11] G. Hörmann and M. Oberguggenberger. Elliptic regularity and solvability for partial differential equations with Colombeau coefficients. *Electron. J. Diff. Eqns.*, 2004(14):1–30, 2004.
- [12] A. E. Hurd and D. H. Sattinger. Questions of existence and uniqueness for hyperbolic equations with discontinuous coefficients. *Trans. Amer. Math. Soc.*, 132:159–174, 1968.
- [13] M. Oberguggenberger. Hyperbolic systems with discontinuous coefficients: examples. In B. Stanković, E. Pap, S. Pilipović, and V. S. Vladimirov, editors, *Generalized Functions, Convergence Structures, and Their Applications*, pages 257–266, New York, 1988. Plenum Press.
- [14] M. Oberguggenberger. *Multiplication of distributions and applications to partial differential equations*. Longman Scientific & Technical, 1992.